

The Essentials of Urine Drug Screens

Drug screenings play a vital role in patient care and can help pharmacists aid in a diagnosis, document adherence and potentially identify diversion of a substance. The tips below provide reminders on how to interpret urine drug screens including the cross reactivity of different substances.



Reasons for Unexpected Results



Positive

- Use of illicit drug or non-prescribed medication
- False positive/cross reactivity
- Contamination




Negative

- Has not used recently (uses PRN, ran out early)
- Test does not detect specific medication (e.g., synthetic opiates)
- High cutoff values
- Fast metabolizer
- Diversion
- Laboratory error
- Tampering of sample or dilution of urine



Expected Metabolites for Opioids with Confirmation Testing

SUBSTANCE	POSSIBLE URINARY METABOLITES ON CONFIRMATION TESTING (e.g., GC/MS)
Morphine	Morphine; Hydromorphone (minor metabolite)
Codeine	Codeine; Morphine; Hydrocodone (minor metabolite)
Hydrocodone	Hydrocodone; Hydromorphone
Hydromorphone	Hydromorphone
Heroin	6- Monoacetylmorphine (6-MAM); Morphine
Oxycodone	Oxycodone; Oxymorphone



Evaluation of Urine

Urine creatine	> 20 mg/dl
Specific gravity	> 1.002
pH	4.5 – 8.0
Temperature	32 - 38 °C (89.6 – 100.4 °F)
Appearance	Yellow, free of bubbles or abnormal smells



Potential Positive Results with Immunoassays and Detection Time in the Urine

SUBSTANCE	POTENTIAL POSITIVES	LENGTH OF TIME DETECTED IN URINE
Amphetamines	l-methamphetamine (Vick's® Inhaler) Amantadine Aripiprazole Atomoxetine Bupropion Chlorpromazine Desipramine Ephedrine Labetalol Methylphenidate MDMA Phentermine Promethazine Pseudoephedrine Phenylephrine Ranitidine Selegiline Thioridazine Trazodone	48 hrs
Benzodiazepines	Oxaprozin Sertraline	Short acting – 3 days Long acting > 30 days
Cocaine	Coca leaf products Topical anesthetics containing cocaine	2-4 days
Opiates	Dextromethorphan Quinine Quinolones Rifampin	2-3 days
Marijuana	Baby wash products Dronabinol Efavirenz NSAIDS Pantoprazole	Single use – 3 days Chronic heavy use > 30 days
Methadone	Diphenhydramine Doxylamine Verapamil	3 days
PCP	Dextromethorphan Diphenhydramine Doxylamine Ibuprofen Imipramine Ketamine Lamotrigine MDPV Meperidine Tramadol Venlafaxine	8 days
Synthetic Cannabinoids	Lamotrigine	3 days


References

Moeller KE, Kissack JC, Atayee, RS, and Lee, KC. Clinical interpretation of urine drug tests: What clinicians need to know about urine drug screens. Mayo Clinic Proceedings. 2017; 92:774-796.
 Cone EJ, Dickerson S, Paul BD, Mitchell JM. Forensic drug testing for opiates. V. Urine testing for heroin, morphine, and codeine with commercial opiate immunoassays. J Anal Toxicol. 1993;17:156-164.
 Verstraete AG. Detection times of drugs of abuse in blood, urine, and oral fluid. Ther Drug Monit. 2004;200-205.

Advance your knowledge on this topic through a complimentary, ACPE approved webinar available at cpnp.org/370286



This toolkit is presented by the College of Psychiatric and Neurologic Pharmacists (CPNP) through a grant supported by the Providers Clinical Support System (PCSS).



Providers Clinical Support System

Funding for this initiative was made possible (in part) by grant no. 5H79T1025595-03 from SAMHSA. The views expressed in written conference materials or publications and by speakers and moderators do not necessarily reflect the official policies of the Department of Health and Human Services; nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government.